

Phenotypic and Molecular Screening of Laccase-producing Bacteria Isolated from Automobile Workshop Soil Samples in Ado-Ekiti

Temitayo Omotunde Olowomofe¹, Olaoluwa Jacob Oluyeye¹, Paul Ikechukwu Orjiakor¹, Ayodele Oluwayemisi Ogunlade², Solomon Temitayo Olaoye¹

¹Department of Microbiology, Faculty of Science, Ekiti State University, Ado-Ekiti, Nigeria

²Department of Food Technology, Federal Polytechnic, Ado-Ekiti, Nigeria

Email address:

motunde21@yahoo.com (T. O. Olowomofe), temitayo.olowomofe@eksu.edu.ng (T. O. Olowomofe)

To cite this article:

Temitayo Omotunde Olowomofe, Olaoluwa Jacob Oluyeye, Paul Ikechukwu Orjiakor, Ayodele Oluwayemisi Ogunlade, Solomon Temitayo Olaoye. Phenotypic and Molecular Screening of Laccase-producing Bacteria Isolated from Automobile Workshop Soil Samples in Ado-Ekiti. *International Journal of Microbiology and Biotechnology*. Vol. 5, No. 3, 2020, pp. 97-102. doi: 10.11648/j.ijmb.20200503.14

Received: April 21, 2020; **Accepted:** May 13, 2020; **Published:** June 20, 2020

Abstract: This work was carried out to isolate hydrocarbon-degrading bacteria from oil contaminated soil sample in Ado-Ekiti and screen them for laccase production. Soil samples were collected and analyzed using standard microbiological techniques. The isolates were initially screened for hydrocarbon degrading ability on minimal salt medium supplemented with 1% crude oil and incubated for 14 days. The isolates were further screened for their ability to produce laccase enzyme using plate screening and molecular techniques. Four of the isolates that gave the best results on tannic-agar plates were selected for PCR amplification of laccase gene using specific primers. The isolates were identified as *Lactobacillus sakei*, *Pseudomonas aeruginosa*, *Bacillus cereus* and *Gracilibacter thermotolerans* based on 16SrRNA sequencing. The DNA of these bacteria amplified the primer specific for laccase gene with 1300bp, 1400bp, 1600bp and 350bp respectively. For bioremediation to be effective, microorganisms must enzymatically attack the pollutants and convert them to harmless products. Therefore, laccase production potentials in these bacteria make them useful in bioremediation as laccase is known to break heavy phenol containing hydrocarbons. Further work can be done to determine the activity of this enzyme during the degradation of crude oil.

Keywords: Bioremediation, Laccase, PCR Amplification, Hydrocarbon-degrading Bacteria

1. Introduction

One of the major environmental problems faced by the world today is the contamination of soil, water, and air by toxic chemicals. Organic and inorganic pollutants from industries and extensive use of pesticides in agriculture have immensely contributed to this problem. Many of these compounds, such as polycyclic aromatic hydrocarbons (PAH), pentachlorophenols (PCP), polychlorinated biphenyls (PCB), 1,1,1-trichloro-2,2-bis(4-chlorophenyl) ethane (DDT), benzene, toluene, ethylbenzene, and xylene (BTEX) as well as trinitrotoluene (TNT), are persistent in the environment and are known to have carcinogenic and/or mutagenic effects [1, 2]. Bioremediation has been the best option for converting these hazardous pollutants into less hazardous compounds by utilizing microorganisms. Hydrocarbon-degrading

microorganisms are widely distributed in the soils, where they play important role in remediating polluted soils. When they come into contact with complex organic materials, extracellular enzymes are released to convert high molecular weight materials into diffusible fractions, which could be transported through the cell wall for assimilation. The role of enzymes in keeping the environment clean is immense; being implemented in the biodegradation of various environmental pollutants and biocatalyzing various reactions by substituting the environmentally hazardous and toxic chemical catalysts offering an environment friendly alternative [3]. Laccases (benzenediol: oxygen oxidoreductases; EC 1.10.3.2) are copper-containing enzymes that belong to the so-called blue copper oxidases. These enzymes are responsible for the oxidation of a variety of phenolic compounds as well as aromatic amines with the reduction of molecular oxygen to water [4]. In addition, in the presence of small molecular-

weight compounds named redox mediators, laccases are able to oxidize non-phenolic structures [5]. This together with the fact that they only use molecular oxygen as a co-substrate instead of hydrogen peroxides as used by peroxidases, make laccases very attractive for different biotechnological applications [6]. Laccases are also useful for the decomposition of azo dyes by oxidative methods [7].

Laccase and laccase-producing microorganisms play an important role in bioremediation of aromatic substances from contaminated soils, industrial pollutants and xenobiotics. They are able to oxidize toxic organic pollutants, such as polycyclic aromatic hydrocarbons and chlorophenols [8]. The ability of laccase producing microorganisms or purified laccases to eliminate a wide range of pollutants is currently one of the most interesting subjects for researchers in environmental biotechnology [9]. Screening of laccase-producing microorganisms is paramount for selecting suitable laccase producing organisms. Screening of microorganisms for enzyme production is usually done by qualitative and quantitative assays which are accompanied with some imprecisions. Hence, molecular screening presents a rapid and more sensitive procedure for determining the enzymatic competence of microorganisms. In this study, hydrocarbon-degrading bacteria were screened for laccase enzyme using plate screening and molecular methods.

2. Materials and Methods

2.1. Collection of Samples

Soil samples were collected in sterile polythene bags at a depth of 10cm from eight different mechanic workshops in Ado-Ekiti. Collected samples were transported to the laboratory for analyses.

2.2. Isolation of Hydrocarbon-utilizing Bacteria

One gram of each sample was introduced into sterile mineral salt medium [NaCl (0.1g/l); MgSO₄·7H₂O (1.2g/l), KH₂PO₄ (1.2g/l), FeSO₄·7H₂O (6.04g/l) distilled water at pH 7.2] supplemented with 1% bonny light crude oil. The set up was incubated at 37°C for 7 days. After 7 days, 1ml of each set up was inoculated into a freshly prepared MSM with 1% of crude oil and incubated at 37°C for another 7 days. After 14 days, ten-fold serial dilution was carried out. Dilutions 10⁵ and 10⁶ were plated using pour plate technique. Plates were incubated at 37°C for 24 hours. Total hydrocarbon-degrading bacteria in the samples were calculated in colony forming units/ml (CFU/ml). Pure cultures were obtained using streak method and stored on agar slants as stock cultures for further analyses.

2.3. Laccase Enzyme Screening

The bacteria isolated were screened for laccase enzyme using tannic acid as a substrate. The isolates were inoculated in Nutrient agar plates containing 5% tannic acid using agar diffusion method [10, 11]. The plates were incubated at room temperature for 24h. Brown-black zones were observed at the

center of some plates and the diameter of the zones was measured. Isolates with positive reaction were selected for molecular screening.

2.4. Quantitative Assay of Laccase

Laccase activity was measured using 2 mM ABTS {2, 2-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid)} as a substrate [12]. The reaction mixture consists of 10 mM sodium acetate buffer (pH 5) containing ABTS at a concentration of 2 mM and 50 to 100 µl cell free supernatant. The absorbance was measured at 470 nm using spectrophotometer (JENWAY6705) [13]. One unit of laccase was defined as the amount of the enzyme required to transform 1 mol substrate per min under standard assay conditions.

2.5. Identification of Isolates

The isolates were identified based on morphological and biochemical characterization. The identities of the best four isolates on plate screening test were confirmed with 16S rRNA sequencing.

2.6. DNA Extraction and PCR Amplification

The bacteria DNA was extracted using Jena Bioscience Kit by following the manufacturer's instructions. The PCR reaction was carried out using the Solis Biodyne 5X HOT FIREPol Blend Master mix. PCR was performed in 25 µl of a reaction mixture, and the reaction concentration was brought down from 5x concentration to 1X concentration containing 1X Blend Master mix buffer (Solis Biodyne), 1.5mM MgCl₂, 200µM of each deoxynucleoside triphosphates (dNTP)(Solis Biodyne), 25pMol of each primer (Jena Bioscience, Germany), 2 unit of Hot FIREPol DNA polymerase (Solis Biodyne), Proofreading Enzyme, 5µl of the extracted DNA, and sterile distilled water was used to make up the reaction mixture. Thermal cycling was conducted in an Peltier thermal cycler (PTC100) (MJ Research Series) for an initial denaturation of 95°C for 15 minutes followed by 35 amplification cycles of 30 seconds at 95°C; 1 minute at 60°C (Lac; (5'ATGACACTTGAAAAATTTGTGGATGCTCTCCCC3' and 5'CTATTTATGGGGATCAGTTATATCCATCGG3'), 65°C (27F; (5'ATGAGTGGCTTGACGCAGGCGCTGCTG3' and 27R (5'CTAGCGCGGGTCCAGCCAGACCAACGATGC 3') and 1 minute 30 Seconds at 72°C. This was followed by a final extension step of 10 minutes at 72°C. The amplification product was separated on a 1.5% agarose gel and electrophoresis was carried out at 80V for 1 hour 30 minutes. After electrophoresis, DNA bands were visualized by ethidium bromide staining. 100bp DNA ladder (Solis Biodyne) was used as DNA molecular weight marker.

2.7. Statistical Analysis

Data obtained from the study were analyzed using Microsoft Excel 2016.

3. Results and Discussion

In automobile workshops, there is a constant change in the soil microorganism as a result of deliberate spillage of used engine oil. These alter the biomass and ecology of the soil such that both microbial communities and grasses can no longer grow on the soil spots. The colour and texture of the soil are affected; this leads to different microbial flora establishment in an attempt to remedy the petroleum product spillage [14]. Laccase as biocatalysts have received lots of attention because of their high capacity of oxidizing phenolic and other aromatic compounds. These advantages make laccases suitable for bioremediation of polluted sites.

Isolation and screening of hydrocarbon-utilizing bacteria:

Table 1. Total Bacteria Count ($\text{Log}_{10}\text{CFU/ml}$) And Total Hydrocarbon-Degrading Bacteria Count ($\text{Log}_{10}\text{CFU/ml}$) Of Soil Samples Obtained from Automobile Workshops in Ado-Ekiti.

SAMPLES	Total bacteria count ($\text{Log}_{10}\text{CFU/ml}$)	Total hydrocarbon- utilizing bacteria (HUB) count ($\text{Log}_{10}\text{CFU/ml}$)	% HUB
S1	7.08 \pm 0.04	5.31 \pm 0.05	75
S2	8.11 \pm 0.02	6.83 \pm 0.01	84
S3	6.16 \pm 0.06	4.64 \pm 0.01	75
S4	7.28 \pm 0.02	4.85 \pm 0.02	66
S5	8.30 \pm 0.05	6.31 \pm 0.01	76
S6	7.41 \pm 0.04	5.46 \pm 0.04	73
S7	8.37 \pm 0.03	4.64 \pm 0.03	55
S8	6.88 \pm 0.04	5.50 \pm 0.05	79
AVERAGE	7.45	5.44	73

Laccase enzyme screening:

Twenty-nine hydrocarbon degraders recovered from the mechanic workshop soil samples showed different reactions on the tannic acid agar. Oxidation of tannic acid was evident by development of black coloration (Figure 1). Four isolates (MW 4, MW 5, MW 8 & MW 11) showed oxidation of tannic-acid within 24h, three isolates (MW 2, MW 15, & MW 21) oxidized tannic acid after 72h and the remaining isolates did not show any coloration. It was observed that incubation period required for proper growth and development of colored precipitate varied with individual organism.

The seven isolates that gave a positive reaction on tannic-acid agar were subjected to further quantitative laccase screening.




Figure 1. Sample plate showing laccase screening on tannic acid agar plate.

Quantitative Assay of Laccase

Laccase activity was determined using ABTS [2, 2-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid)] as a substrate over a period of 96h. All the seven isolates that gave positive reaction

Table 1 reports the total bacteria count and total hydrocarbon utilizing bacteria of soil samples from eight automobile workshops. Average count of 7.45 $\text{Log}_{10}\text{CFU/ml}$ and 5.44 $\text{Log}_{10}\text{CFU/ml}$ was recorded respectively. However high percentage of hydrocarbon utilizers were recovered from samples in these locations. Prevalence of hydrocarbon degraders in oil polluted environments have been documented in previous researches [15]. The higher count of HUB could also be attributed to long term exposure of these bacteria to hydrocarbons thereby acclimatized to the environment. This agrees with the work of (Rahman *et al.* [16] who reported that the population levels of hydrocarbon utilizers within the contaminated environment appear to be a sensitive index of environmental exposure to hydrocarbons.

on the tannic-acid agar were subjected to quantitative screening. Laccase activity increased gradually with time to attain the optimum between 48 and 72h and later declined after 96h as shown in Figure 2. Incidentally, isolates MW15 and MW21 which gave late positive reaction (after 72h) on plate screening had the highest laccase activity of 3.43U/ml and 3.09U/ml respectively after 72h while their counterparts MW5 and MW11 which gave fast positive reaction on tannic-acid agar plate had considerably lower activity of 1.56 U/ml and 1.15 U/ml respectively. However, it has been reported that plate assay test for screening based on colour indicator compounds viz. guaiacol and tannic acid are efficient substrates to obtain novel laccase producers [17] but some of the microbes that showed rapid positive reactions on tannic-acid produced low laccase in liquid cultures. This shows that quantification assay is more sensitive than the plate screening method.




Figure 2. Qualitative assay of Laccase in hydrocarbon-degrading bacteria.

Identification of the isolates: The four isolates with the

highest laccase activity were identified as *Lactobacillus sakei*, *Pseudomonas aeruginosa*, *Bacillus cereus* and *Gracilibacter thermotolerans* based on 16SrRNA sequencing.

Sample A- *Lactobacillus sakei* strain DSM 20017 16S ribosomal RNA gene, partial sequence

Sequence ID: gi|343201717|NR_042443.1Length: 1561Number of Matches: 1
Related Information
Range 1: 257 to 291GenBankGraphics Next Match Previous Match

Table 2. % Sequence similarities of Isolate M4 with closest relatives from the Genbank database.

Alignment statistics for match #1				
Score	Expect	Identities	Gaps	Strand
46.4 bits (50)	2e-04	31/35 (89%)	0/35 (0%)	Plus/Plus
Query	209	CGCGGTCCATAAGTTATTTGGGGAGGTAAAGGCTC	243	
Sbjct	257	CGCGGTGCATTAGTTAGTTGGTGAGGTAAAGGCTC	291	

Sample B-*Pseudomonas aeruginosa* strain DSM 50071 16S ribosomal RNA, complete sequence

Sequence ID: gi|219846486|NR_026078.1Length: 1537Number of Matches: 1

Related Information

Range 1: 10 to 934GenBankGraphics Next Match Previous Match

Table 3. % Sequence similarities of Isolate M8 with closest relatives from the Genbank database.

Alignment statistics for match #1				
Score	Expect	Identities	Gaps	Strand
1548 bits (1716)	0.0	906/929 (98%)	9/929 (0%)	Plus/Plus
Query	1	AGTTTGATCCTGGCTCAGATTGAACGCTGGCGGCAGG--CCT--AACACATGCAAGTCGA	56	
Sbjct	10	AGTTTGATCATGGCTCAGATTGAACGCTGGCAGCAGGGGCTTCAACACATGCAAGTCGA	69	
Query	57	GCGGATGAAGGGAGCTTGCTCCTGGATTACGCGGCGGACGGGTGAGTAATGCCTAGGAAT	116	
Sbjct	70	GCTTATGAAGGGAGCTTGC-CTTGGATTACGCGGCGGACGGGTGAGTAATGCCTAGGAAT	128	
Query	117	CTGCCTGGTAGTGGGGGATAACGTCCGGAACGGCGCTAATACCGCATACTGCTGAGG	176	
Sbjct	129	CTGCCTGGTAGTGGGGGATAACGTCCGGAACGGCGCTAATACCGCATACTGCTGAGG	188	
Query	177	GAGAAAGTGGGGGATCTTCGGACCTCACGCTATCAGATGAGCCTAGGTGCGATTAGCTAG	236	

Sample C -*Bacillus cereus* ATCC 14579 16S ribosomal RNA (rrnA) gene, complete sequence

Sequence ID: gi|444304116|NR_074540.1Length: 1512Number of Matches: 2

Related Information

Range 1: 134 to 629GenBankGraphics Next Match Previous Match

Table 4. % Sequence similarities of Isolate M21 with closest relatives from the Genbank database.

Alignment statistics for match #1				
Score	Expect	Identities	Gaps	Strand
666 bits (738)	0.0	447/496 (92%)	2/496 (0%)	Plus/Plus
Query	102	CCCATAAGATTGGATC-CTCCGGAACCGGGGCTAAAACGGAAAAACATTTTGAACCG	160	
Sbjct	134	CCCATAAGACTGGGATAACTCCGGGAACCGGGGCTAATACCGGATAACATTTTGAACCG	193	
Query	161	CAGGGTTCAAAATTGAAAGGCGGCTTCGGCTGTCACTTATGGATGGACCCGCGTCGCATT	220	
Sbjct	194	CATGGTTCGAAATTGAAAGGCGGCTTCGGCTGTCACTTATGGATGGACCCGCGTCGCATT	253	
Query	221	AGCTAGTTGGTGAGGTAACGGCTCACCAGGGCAACAATGCGTACCCAACCTGAGAGGGTG	280	
Sbjct	254	AGCTAGTTGGTGAGGTAACGGCTCACCAGGGCAACGATGCGTAGCCGACCTGAGAGGGTG	313	
Query	281	ATCGGCCACCTGGGACTGAAACACGGCCCAACTCCTACGGGAGGCAGCAGTAGGGAAT	340	
Sbjct	314	ATCGGCCACACTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTAGGGAAT	373	
Query	341	CTTCGCAATGGACAAAAGTCTGACGGACACCCCGCGTGAGTGATGAAGGTTTTCGGG	400	
Sbjct	374	CTTCGCAATGGACAAAAGTCTGACGGAGCAACCCCGCGTGAGTGATGAAGGCTTTCGGG	433	
Query	401	TCGTAAACTCTGTTGTTGGGGAAAAACAAGTGCTAGTTGAATAAGCTGGCCCCCTTACG	460	
Sbjct	434	TCGTAAACTCTGTTGTTAGGGAAGAACAAAGTGCTAGTTGAATAAGCTGGCACCTTGACG	493	

Sample D-*Gracilibacter thermotolerans* strain JW/YJL-S1 16S ribosomal RNA gene, partial sequence
Sequence ID: gi|636559633|NR_115693.1 Length: 1526 Number of Matches: 1
Related Information
Range 1: 1383 to 1416 GenBank Graphics Next Match Previous Match

Alignment statistics for match #1				
Score	Expect	Identities	Gaps	Strand
31.9 bits (34)	3.0	28/34 (82%)	1/34 (2%)	Plus/Plus

27F @ 65

Key: M: 100 bp DNA ladder, Invitrogen (positive control), -ve: negative control line. Lane A=*Lactobacillus sakei*, Lane B=*Pseudomonas aeruginosa*, Lane C=*Bacillus cereus* & Lane D=*Gracilibacter thermotolerans*.

- [1] Cerniglia, N. and Sutherland, M. (2001). *Application Environmental Microbiology*. 63: 4120-4122.
- [2] Clemente, A., Dimcheva, N., Ferapontova, E. E., Gorton, L., Ruzgas, T., Stoica, L., Shleev, S., Yaropolov, A. I., Haltrich, D., Thorneley, R. N. and Austin, S. D. (2001). *Application of Microbiology and Biotechnology*. 16: 1074-1092.
- [3] Bird, C. E. Cavanaugh, E. G. Dickerman, J. C. and Fernandes, S. R. (2008). A Study of Vapor Control Methods for Gasoline Marketing Operations. *U.S. Environmental Protection Agency*. 171: 31-36.

- [4] Zouari-Mechichi, H., Mechichi, T., Dhoub, A., Sayadi, S., Martinez, A. T., mMartinez, M. J., (2005). Laccase purification and characterization from *Trametes trogii* isolated in Tunisia: decolorization of textile dyes by the purified enzyme. *Enzyme Microbial Tech.*, 39: 141-148.
- [5] Desai, S., Nityanand, C. (2011). Microbial laccases and their applications: a review. *Asian J Biotechnol*, 3 (2): 98-124.
- [6] Bains J, Capalash N, Sharma P (2003). Laccase from a non-melanogenic, alkalotolerant-proteobacterium JB isolated from industrial wastewater drained soil. *Biotechnol Lett* 25 (1): 155-1159.
- [7] Michael, M. T., M. G. Georg and R. Astrid (2005). Degradation of Azo dyes by lacase and ultrasound treatment. *Apl. Environ. Microbiol.* 71: 260-2607.
- [8] Elsayed, M. A., Hassan, M. M., Elshafei, A. M., Haroun, B. M. Ot man, A. M. (2012). Optimization of cultural and nutritional parameters for the production of laccase by *Pleurotus ostreatus* ARC280. *British Biotechnology Journal*, 2 (3), 115-132.
- [9] Baldrian, P. (2006). "Fungal laccases-occurrence and properties," *FEMS Microbiology Reviews*, vol. 30, no. 2, pp. 215-242.
- [10] Sivakumar R, Rajendran R, Balakumar C, Tamilvendan M (2010). Isolation, screening and optimization of production medium for thermostable laccase production from *Ganoderma* sp. *Int. J. Eng. Sci. Technol.* 2 (12): 7133-7141.
- [11] Pointing, S. B. (1999). Qualitative methods for the determination of lignocellulolytic enzyme production by tropical fungi. *Fungal Diversity* 2, 17-33.
- [12] Bourbonnais, R., Leech, D. and Paice, M. G. (1998). "Electrochemical analysis of the interactions of laccase mediators with lignin model compounds," *Biochimica et Biophysica Acta*, vol. 1379, no. 3, pp. 381-390.
- [13] Majcherczyk, A., Johannes, C. and A. Huttermann, A. (1998). "Oxidation of polycyclic aromatic hydrocarbons (PAH) by laccase of *Trametes versicolor*," *Enzyme and Microbial Technology*, vol. 22, no. 5, pp. 335-341.
- [14] Barathi, S. and Vasudevan, N. (2001). Utilization of petroleum hydrocarbons by *pseudomonas fluorescens* isolated from petroleum contaminated soil. *Environmental International*. 26: 413-416.
- [15] Olowomofe T. O, Oluyeye J. O, Sowole D. O (2017). Isolation, screening and characterization of hydrocarbon-utilizing bacteria isolated from bitumen-contaminated surface water in Agbabu, Ondo State. *J Adv Biol Biotechnol* 15: 1-9.
- [16] Kiiskinen, L. L., Rättö, M., Kruus, K. (2004). Screening for novel laccase-producing microbes. *Journal of applied microbiology*, 97 (3): 640-646.
- [17] Rahman KSM, Rahman TJ, Kourkoutas Y, Petsas I, Marchant R, Banat I. M. (2003). Enhanced bioremediation of N-alkane in petroleum sludge using bacterial consortium amended with rhamnolipid and micronutrients. *Biores. Technol.*, 90: 159-168.